Gammopathies Monoclonales

Place de l'analyse des immunoglobulines monoclonales

*Dr Yannick CHANTRAN, PharmD, PhD,
Chef du service d'Immunologie Biologique, Sorbonne Université
Hôpital Saint-Antoine*

Myélome - Epidémiologie

- 1.8 % des nouveaux cas de cancers
- 2.1 % des décès par cancers
- 2^e hémopathie maligne
- ↑ Incidence
- ↓ Mortalité
- Sex ratio H:F 1.5:1
- Age médian au diagnostic 69 ans

Myélome: Physiopathologie et signes cliniques

Du MGUS au Myélome: un continuum

Figure 2. Probability of Progression to Active Multiple Myeloma or Primary Amyloidosis in Patients with Smoldering Multiple Myeloma or Monoclonal Gammopathy of Undetermined Significance (MGUS). I bars denote 95% confidence intervals.

Immunoglobuline monoclonale & Diagnostic du MGUS

- EPS
- IF
- EPU ou IFU
- FLC

Disorder	Disease Definition
Non-IgM MGUS	All three criteria must be met:
	Serum monoclonal protein (non-IgM type) < 3 g/dL
	Clonal bone marrow plasma cells < 10%*
	Absence of end-organ damage such as CRAB features that can be attributed to the plasma cell proliferative disorder
Light-Chain MGUS	All criteria must be met:
	Abnormal FLC ratio (< 0.26 or > 1.65)
	Increased level of the appropriate involved light chain (increased kappa FLC in patients with ratio > 1.65 and increased lambda FLC in patients with ratio < 0.26)
	No immunoglobulin heavy-chain expression on immunofixation
	Absence of end-organ damage that can be attributed to the plasma cell proliferative disorder
	Clonal bone marrow plasma cells < 10%
IgM MGUS	Urinary monoclonal protein < 500 mg/24 h
	All three criteria must be met:
	Serum IgM monoclonal protein < 3 gm/dL
	Bone marrow lymphoplasmacytic infiltration < 10%
	No evidence of anemia, constitutional symptoms, hyperviscosity, lymphadenopathy, or hepatosplenomegaly that can be attributed to the underlying lymphoproliferative disorder.

Immunoglobuline monoclonale & Pronostic des MGUS

Prognostic factor	Hazard ratio (95% confidence Interval)*
Abnormal FLC ratio	2.6 (1.7-4.2)
Serum M protein size > 15 g/L	2.4 (1.7-3.5)
IgA, IgM, or biconal IgA plus IgM	2.6 (1.7-4.0)

- EPS
- IF
- EPU ou IFU
- FLC

* $P < .001$ for each variable.

Immunoglobuline monoclonale & Suivi des MGUS

*Evolving MGUS is defined as an increase of M-protein of at least 10% by the third year, confirmed by two consecutive measurements separated by at least 1 month.

Immunoglobuline monoclonale & Diagnostic des SMM & MM

	EPS
	IF
	EPU ou IFU
	FLC

Smoldering MM	<p>Both criteria must be met:</p> <p style="background-color: #f4a460;">Serum monoclonal protein (IgG or IgA) $\geq$ 3 gm/dL, or urinary monoclonal protein $\geq$ 500 mg per 24 h and/or</p> <p style="background-color: #ffff00;">clonal bone marrow plasma cells 10%–60%</p> <p>Absence of myeloma-defining events or amyloidosis</p>
MM	<p>Both criteria must be met:</p> <p>Clonal bone marrow plasma cells $\geq$ 10% or biopsy-proven bony or extramedullary plasmacytoma</p> <p>Any one or more of the following myeloma defining events:</p> <p>Evidence of end organ damage that can be attributed to the underlying plasma cell proliferative disorder, specifically:</p> <p style="background-color: #f4a460;">Hypercalcemia: serum calcium $>$ 0.25 mmol/L ($>$ 1 mg/dL) higher than the upper limit of normal or $>$ 2.75 mmol/L ($>$ 11 mg/dL)</p> <p style="background-color: #f4a460;">Renal insufficiency: creatinine clearance $<$ 40 mL/min or serum creatinine $>$ 177 μmol/L ($>$ 2 mg/dL)</p> <p style="background-color: #f4a460;">Anemia: hemoglobin value of $>$ 2 g/dL below the lower limit of normal, or a hemoglobin value $<$ 10 g/dL</p> <p style="background-color: #f4a460;">Bone lesions: one or more osteolytic lesions on skeletal radiography, CT, or PET-CT</p> <p>Clonal bone marrow plasma cell percentage $\geq$ 60%</p> <p style="background-color: #a6c9e0;">Involved: uninvolved serum FLC ratio $\geq$ 100 (involved FLC level must be $\geq$ 100 mg/L)</p> <p style="background-color: #a6c9e0;">$>$ 1 focal lesion on MRI studies (at least 5 mm in size)</p>

Immunoglobuline monoclonale & Pronostic du SMM

- EPS
- IF
- EPU ou IFU
- FLC

Figure 3. Probability of Progression to Active Multiple Myeloma or Primary Amyloidosis in Patients with Smoldering Multiple Myeloma among Three Risk Groups.

The proportion of plasma cells in bone marrow and the serum monoclonal protein level were combined to create a risk-stratification model with three distinct prognostic groups. At 15 years, the cumulative probability of progression was 87% for the 106 patients in group 1 (bone marrow plasma cells, $\geq 10\%$; monoclonal protein level, ≥ 3 g per deciliter), 70% for the 142 patients in group 2 (plasma cells, $\geq 10\%$; monoclonal protein level, < 3 g per deciliter), and 39% for the 27 patients in group 3 (plasma cells, $< 10\%$; monoclonal protein level, ≥ 3 g per deciliter).

Immunoglobuline monoclonale & Pronostic du SMM

- EPS
- IF
- EPU ou IFU
- FLC

*Evolving MGUS is defined as an increase of M-protein of at least 10% by the third year, confirmed by two consecutive measurements separated by at least 1 month.

Immunoglobuline monoclonale & Pronostic du MM

- EPS
- IF
- EPU ou IFU
- FLC

TABLE 4 Revised international staging system for myeloma⁵⁴

Stage
<p>Stage 1</p> <p>All of the following:</p> <ul style="list-style-type: none"> • Serum albumin ≥ 3.5 gm/dL • Serum beta-2-microglobulin < 3.5 mg/L • No high-risk cytogenetics • Normal serum lactate dehydrogenase level
<p>Stage II</p> <ul style="list-style-type: none"> • Not fitting Stage I or III
<p>Stage III</p> <p>Both of the following:</p> <ul style="list-style-type: none"> • Serum beta-2-microglobulin > 5.5 mg/L • High-risk cytogenetics [t(4;14), t(14;16), or del(17p)] or Elevated serum lactate dehydrogenase level

?

Immunoglobuline monoclonale & Pronostic du MM

- EPS
- IF
- EPU ou IFU
- FLC

TABLE 4 Revised international staging system for myeloma⁵⁴

Stage
<p>Stage 1</p> <p>All of the following:</p> <ul style="list-style-type: none"> • Serum albumin ≥ 3.5 gm/dL • Serum beta-2-microglobulin < 3.5 mg/L • No high-risk cytogenetics • Normal serum lactate dehydrogenase level
<p>Stage II</p> <ul style="list-style-type: none"> • Not fitting Stage I or III
<p>Stage III</p> <p>Both of the following:</p> <ul style="list-style-type: none"> • Serum beta-2-microglobulin > 5.5 mg/L • High-risk cytogenetics [t(4;14), t(14;16), or del(17p)] or Elevated serum lactate dehydrogenase level

?

IgA MM	<ul style="list-style-type: none"> ↑ ISS-3 ↑ anemie ↑ t(4;14)
FLC MM	<ul style="list-style-type: none"> ↑ IR ↑ t(11;14)

Immunoglobuline monoclonale & Réponse au traitement

Standard IMWG response criteria

Stringent complete response	Complete response as defined below plus normal FLC ratio** and absence of clonal cells in bone marrow biopsy by immunohistochemistry (κ/λ ratio $\leq 4:1$ or $\geq 1:2$ for κ and λ patients, respectively, after counting ≥ 100 plasma cells)††
Complete response	Negative immunofixation on the serum and urine and disappearance of any soft tissue plasmacytomas and $< 5\%$ plasma cells in bone marrow aspirates
Very good partial response	Serum and urine M-protein detectable by immunofixation but not on electrophoresis or $\geq 90\%$ reduction in serum M-protein plus urine M-protein level < 100 mg per 24 h
Partial response	$\geq 50\%$ reduction of serum M-protein plus reduction in 24 h urinary M-protein by $\geq 90\%$ or to < 200 mg per 24 h;
	If the serum and urine M-protein are unmeasurable, a $\geq 50\%$ decrease in the difference between involved and uninvolved FLC levels is required in place of the M-protein criteria;
Minimal response	If serum and urine M-protein are unmeasurable, and serum-free light assay is also unmeasurable, $\geq 50\%$ reduction in plasma cells is required in place of M-protein, provided baseline bone marrow plasma-cell percentage was $\geq 30\%$. In addition to these criteria, if present at baseline, a $\geq 50\%$ reduction in the size (SPD)§§ of soft tissue plasmacytomas is also required
	$\geq 25\%$ but $\leq 49\%$ reduction of serum M-protein and reduction in 24-h urine M-protein by 50–89%. In addition to the above listed criteria, if present at baseline, a $\geq 50\%$ reduction in the size (SPD)§§ of soft tissue plasmacytomas is also required
Stable disease	Not recommended for use as an indicator of response; stability of disease is best described by providing the time-to-progression estimates. Not meeting criteria for complete response, very good partial response, partial response, minimal response, or progressive disease
Progressive disease ¶¶,	Any one or more of the following criteria:
	Increase of 25% from lowest confirmed response value in one or more of the following criteria:
	Serum M-protein (absolute increase must be ≥ 0.5 g/dL);
	Serum M-protein increase ≥ 1 g/dL, if the lowest M component was ≥ 5 g/dL;
	Urine M-protein (absolute increase must be ≥ 200 mg/24 h);
In patients without measurable serum and urine M-protein levels, the difference between involved and uninvolved FLC levels (absolute increase must be > 10 mg/dL);	
In patients without measurable serum and urine M-protein levels and without measurable involved FLC levels, bone marrow plasma-cell percentage irrespective of baseline status (absolute increase must be $\geq 10\%$);	
Appearance of a new lesion(s), $\geq 50\%$ increase from nadir in SPD§§ of > 1 lesion, or $\geq 50\%$ increase in the longest diameter of a previous lesion > 1 cm in short axis;	
$\geq 50\%$ increase in circulating plasma cells (minimum of 200 cells per μL) if this is the only measure of disease

Quelle place pour l'électrophorèse et l'immunofixation des urines ?

- Ne fait pas partie des critères diagnostiques du MM
- **Mais** fait partie des critères pour différencier FLC SMM Vs FLC MGUS
- **Ou** si suspicion d'amylose AL

- Préférentiellement sur des urines de 24h
- Préférentiellement sur des urines concentrées (Levinson, 2011)

- Exclusion d'une gammopathie dans le diagnostic d'une pathologie rénale
- Evaluation de l'atteinte rénale dans le bilan d'une gammopathie

Hors du continuum: les Gammopathies Monoclonales de Signification Clinique

Table 1: Prevalence of monoclonal gammopathies in the Mayo Clinic practice,

Monoclonal gammopathies: Mayo Clinic 1960–2008					
Disease	Number of cases	%	Classification		
			Premalignant	Malignant	Protein/low tumor burden
MGUS	23,179	58.0	X		
Multiple myeloma	6,974	17.5		X	
Primary amyloidosis	3,781	9.5			X
SMM	1,494	3.7	X		
Lymphoproliferative disease	1,298	3.3		X	
Macroglobulinemia	940	2.4		X	
Plasmacytoma	774	1.9		X	
Bence Jones proteinuria	450	1.1	X		
Cryoglobulinemia	379	0.9			X
POEMS syndrome	217	0.5			X
Light chain deposition disease	113	0.3			X
Plasma cell leukemia	90	0.2		X	
Cold agglutinin disease	74	0.2			X
Acquired Fanconi syndrome	43	0.1			X
Scleromyxedema	31	0.1			X
BHPW	31	0.1			X
Heavy-chain disease	31	0.1		X	
Capillary leak syndrome	29	0.1			X
Total	39,929	100			

MGCS
due à une Igmo
pathogène

MGUS, monoclonal gammopathy of undetermined significance; SMM, smoldering multiple myeloma; POEMS, polyneuropathy, organomegaly, endocrinopathy, monoclonal protein, skin changes; BHPW, benign hypergammaglobulinemia purpura of Waldenström's. Adapted from reference [1].

Atteintes d'organes et diagnostic des MGCS

Particularités immunochimiques des MGCS : l'exemple des cryoglobulinémies

Case 1

Case 1

Case 2

Case 2

Particularités immunochimiques des MGCS : l'exemple du scléromyxoedème

Patient	Gender/Age	Severity	Monoclonal Ig isotype	Electrophoretic mobility (normalized value)
1	H/60	Non severe	IgG1 λ	Medium/cathodic (0.98)
2	F/29	Severe, DNS, cardiomyopathy	IgG1 λ	Cathodic (1.00)
3	H/75	Non severe, cutis laxa-like eruption, MBL	IgG1 κ	Cathodic (1.02)
4	H/73	Non severe	IgG1 λ	Cathodic (1.01)
5	F/71	Non severe	IgG1 λ	Cathodic (1.02)
6	F/29	Severe, DNS	IgG1 κ	Medium (0.92)
7	F/75	Non severe	IgG1 λ	Cathodic (1.00)
8	F/69	Non severe	IgG1 κ	Cathodic (1.10)
9	M/51	Non severe	IgG1 λ	Medium (0.89)
10	M/53	Non severe	IgG1 κ	Anodic (0.86)
11	M/55	Severe, recurrent DNS	IgG1 κ	Cathodic (1.11)
12	F/75	Non severe	IgG1 λ	Cathodic (1.05)

Physiopathologie & Implications pour la prise en charge des MGCS

frontiers | Frontiers in Immunology

Monoclonal gammopathies of clinical significance (MGCS): In pursuit of optimal treatment

Artem Oganesyan^{1,2*}, Andrew Gregory³, Florent Malard⁴,
Nerses Ghahramanyan², Mohamad Mohty⁴,
Dickran Kazandjian⁵, Arsène Mekinian^{6,7},
and Yervand Hakobyan^{1,2}, on behalf of MINHEMON

TYPE Review

PUBLISHED 23 November 2022

DOI 10.3389/fimmu.2022.1045002

TABLE 1 Potential mechanisms of MGCS-related diseases.

Proposed mechanism	MGCS-related diseases
Immunoglobulin deposits	Crystal storing histiocytosis Crystalline keratopathy Maculopathy of monoclonal gammopathy
Autoantibodies	Acquired C1 inhibitor deficiency Acquired von Willebrand disease Xanthomatosis CANOMAD CIDP DADS-M
Uncertain	Clarkson's disease TEMPI syndrome Neutrophilic dermatoses SLOMN Acquired cutis laxa Scleromyxedema Schnitzler syndrome

Ouverture : Analyse des gammopathies par Spectrométrie de Masse

Merci de votre attention !